( PAR	FB 1 5 2	V 3C48 32
-6	& RADEMAP	K QEL

LIST OF PRIOR ART CITED BY APPLICANT (Use several sheets if necessary)

Sheet _	•	•	•
SOPPI		of .	•
UM-00-1		O1	

ATTY. DOCKET NO. 5627\*5

SERIAL NO. 10/018103

APPLICANT:

A. James Mixson

FILING DATE: 11/5/01

GROUP

## U.S. PATENT DOCUMENTS

EXAMINER INITIAL		DOCUMENT NUMBER	DATE	NAME	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
	**	5,354,844	11/11/94	Beug et al.			
	AB	5,554,388	9/10/96	Illum			
	AC	5,736,392	4/7/98	Hawley-Nelson et al.		£	
Í	AD	5,845,435	1/5/99	Bazile et al.			
	AE	5,985,354	11/16/99	Mathiowitz et al.		•	
V	AF	6,051,429	4/18/00	Hawley-Nelson et al.			

## FOREIGN PATENT DOCUMENTS

		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATIO YES	ON NO
N	AG	EP 0 727 223 A1	4/6/95	EPO				
(	, AH	WO 98/22610	5/j22/98	PCT		/		
,	AI	WO 99/42091	8/26/99	PCT				
	N	WO 00/32764	6/8/00	PCT				

## OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages, Etc.)

	AK	al., Membrane Permeabilization and Efficient Gene Transfer by a Peptide Containing Several. Bioconjug Chem 98, 9, 260-267.
1	AL	al., Efficient Gene Transfer by Histidylated Polylysine/pDNA Complexes. <i>Bioconjugate</i> May-Jun; 10(3):4-6-411.
	AMI	, Co-polymer of histidine and lysine markedly enhances transfection efficiency of liposomes. 2000 Oct; 7(19):1698-1705.
	AN	Branched co-polymers of histidine and lysine are efficient carriers of plasmids. <i>Nucleic</i> 2001 Mar 15; 29(6):1334-1340.
The second secon	AO	1., Histidylated oligolysines increase the transmembrane passage and the biological activity of ligonucleotides. Nucleic Acids Res 2000 Jan 15; 28(2):504-512.
	AP	al., Polymer-based gene delivery with low cytotoxicity by a unique balance of side-chain roc Natl Acad Sci USA Jan 30; 98(3):1200-1205.

DATE CONSIDERED

**EXAMINER**